

MIT Open Access Articles

First observation of the rare $B^{+} \rightarrow D^{+}K^{+}\pi^{-}$ decay

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Aaij, R., C. Abellán Beteta, B. Adeva, M. Adinolfi, A. Affolder, Z. Ajaltouni, S. Akar, et al. "First Observation of the Rare $B^{+} \rightarrow D^{+}K^{+}\pi^{-}$ Decay." Phys. Rev. D 93, no. 5 (March 3, 2016). © 2016 CERN, for the LHCb Collaboration

As Published: <http://dx.doi.org/10.1103/PhysRevD.93.051101>

Publisher: American Physical Society

Persistent URL: <http://hdl.handle.net/1721.1/101604>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of use: Creative Commons Attribution



First observation of the rare $B^+ \rightarrow D^+K^+\pi^-$ decayR. Aaij *et al.**

(LHCb Collaboration)

(Received 9 December 2015; published 3 March 2016)

The $B^+ \rightarrow D^+K^+\pi^-$ decay is observed in a data sample corresponding to 3.0 fb^{-1} of pp collision data recorded by the LHCb experiment during 2011 and 2012. The signal significance is 8σ and the branching fraction is measured to be $\mathcal{B}(B^+ \rightarrow D^+K^+\pi^-) = (5.31 \pm 0.90 \pm 0.48 \pm 0.35) \times 10^{-6}$, where the uncertainties are statistical, systematic and due to the normalization mode $B^+ \rightarrow D^-K^+\pi^+$, respectively. The Dalitz plot appears to be dominated by broad structures. Angular distributions are exploited to search for quasi-two-body contributions from $B^+ \rightarrow D_2^*(2460)^0K^+$ and $B^+ \rightarrow D^+K^*(892)^0$ decays. No significant signals are observed and upper limits are set on their branching fractions.

DOI: 10.1103/PhysRevD.93.051101

A key goal of flavor physics is to determine precisely the angle γ of the unitarity triangle constructed from pairs of elements of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [1,2]. The value of $\gamma \equiv \arg[-V_{ud}V_{ub}^*/(V_{cd}V_{cb}^*)]$ is currently known to a precision of only about 10° [3–5], which limits the sensitivity of tests of the Standard Model through global fits to the CKM matrix parameters.

A powerful method to determine γ , known as the GLW method [6,7], is to use $B^+ \rightarrow DK^+$ decays with the neutral D meson decaying to CP eigenstates. The $\bar{b} \rightarrow \bar{c}$ and $\bar{b} \rightarrow \bar{u}$ amplitudes both contribute to the decay, and the sensitivity to γ comes from their interference. A challenge with this method is that the ratio of magnitudes of the suppressed and favored amplitudes, r_B , is not known independently and must be determined simultaneously with γ . This is usually addressed by using in addition other decays of the D meson that provide complementary information on r_B and γ [8,9].

In the case of $B^+ \rightarrow D^{**}K^+$ decays, where D^{**} represents an excited D or \bar{D} meson such as the $D_2^*(2460)$ state which can decay to both $D^\pm\pi^\mp$ and $D\pi^0$, it is possible to obtain a clean determination of r_B [10]. The relative branching fractions of the $\bar{b} \rightarrow \bar{u}$ mediated $B^+ \rightarrow D^{**0}K^+ \rightarrow D^+\pi^-K^+$ and the $\bar{b} \rightarrow \bar{c}$ mediated $B^+ \rightarrow \bar{D}^{**0}K^+ \rightarrow D^-\pi^+K^+$ processes give the value of r_B^2 , while the $B^+ \rightarrow D^{**}K^+ \rightarrow D\pi^0K^+$ final state, where the D meson is reconstructed using CP eigenstate decay modes, provides sensitivity to γ . Decay diagrams for $B^+ \rightarrow \bar{D}_2^*(2460)^0K^+$ and $B^+ \rightarrow D_2^*(2460)^0K^+$ decays are shown in Figs. 1(a) and 1(b).

Knowledge of the resonant structure of $B^+ \rightarrow D^+K^+\pi^-$ and $B^+ \rightarrow D^-K^+\pi^+$ decays is therefore needed. The latter channel has recently been studied with a Dalitz plot analysis [11]. Such a study would be difficult with the low yields expected for $B^+ \rightarrow D^+K^+\pi^-$ decays in the available data samples, but an alternative approach exploiting the angular decay information to separate different spin states is viable in the region of the narrow $D_2^*(2460)^0$ resonance. The same method can also be used to search for $B^+ \rightarrow D^+K^*(892)^0$ decays, which contribute to the $D^+K^+\pi^-$ final state and are of interest since they are mediated by annihilation amplitudes, as shown in Fig. 1(c). A previous LHCb analysis of this mode set an upper limit $\mathcal{B}(B^+ \rightarrow D^+K^*(892)^0) < 1.8 \times 10^{-6}$ at the 90% credibility level [12].

In this paper, the $B^+ \rightarrow D^+K^+\pi^-$ channel is studied for the first time, and searches for $B^+ \rightarrow D_2^*(2460)^0K^+$ and $B^+ \rightarrow D^+K^*(892)^0$ decays are performed. The D^+ meson is reconstructed in the $K^-\pi^+\pi^+$ final state. (The inclusion of charge conjugate processes is implied.) The $B^+ \rightarrow D^-K^+\pi^+$ decay [11] is used for normalization. The analysis is based on 3.0 fb^{-1} of pp collision data collected with the LHCb detector during 2011 and 2012. The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, described in detail in Refs. [13,14]. Simulated events are produced using the software described in Refs. [15–20]. To reduce the risk of biasing results, all analysis procedures were established before the data in the signal region were examined.

Candidates consistent with the decay chains $B^+ \rightarrow D^+K^+\pi^-$ and $B^+ \rightarrow D^-K^+\pi^+$ with $D^\pm \rightarrow K^\mp\pi^\pm\pi^\pm$ are selected. The criteria for $B^+ \rightarrow D^-K^+\pi^+$ and $B^+ \rightarrow D^+K^+\pi^-$ candidates are identical, except for charge requirements, and are very similar to those described in Ref. [11]. A loose preselection is applied before two neural network classifiers (NNs) [21] are used to separate signal decays from background events. The first NN separates true $D^\pm \rightarrow K^\mp\pi^\pm\pi^\pm$ decays from random combinations, and the second (NN2) identifies signal B^+ decays. Both NNs

*Full author list given at the end of the article.




FIG. 1. Decay diagrams for (a) $B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+$, (b) $B^+ \rightarrow D_2^*(2460)^0 K^+$ and (c) $B^+ \rightarrow D^+ K^*(892)^0$ channels.

are trained with a sample of candidates from the topologically similar $B^+ \rightarrow D^- \pi^+ \pi^+$ decay. Additional selection requirements are imposed to reject contributions from specific decay modes. Only candidates with DK mass, $m(DK)$, less than $5140 \text{ MeV}/c^2$ are kept, in order to remove backgrounds from $B^0 \rightarrow D^- K^+$ decays combined with a random pion candidate. Similarly, potential $B^0 \rightarrow D^- \pi^+$ background is eliminated by requiring $m(D\pi) < 4790 \text{ MeV}/c^2$. Contributions from $B^+ \rightarrow D^+ \bar{D}^0$ decays, with $\bar{D}^0 \rightarrow K^+ \pi^-$, are removed by rejecting candidates within $\sim 3\sigma_{K\pi}$ of the \bar{D}^0 mass, where $\sigma_{K\pi}$ is the $K^+ \pi^-$ mass resolution, corresponding to $1830 < m(K\pi) < 1890 \text{ MeV}/c^2$. Although each of these backgrounds affects only one of the final states, the vetoes are applied to both to avoid biasing the relative efficiency.

Signal candidates with invariant mass in the range $5100 - 5800 \text{ MeV}/c^2$ are retained for further analysis. Following all selection requirements, fewer than 1% of events contain more than one candidate; all are retained.

Extended maximum likelihood fits to the distributions of candidates in NN2 output and in B candidate mass are used to determine the yields of $B^+ \rightarrow D^+ K^+ \pi^-$ and $B^+ \rightarrow D^- K^+ \pi^+$ decays. Similar fitting techniques have been used successfully in several previous LHCb analyses [22–25]. A loose requirement is placed on the output of NN2 and the remaining data in each sample are divided into six bins of the NN2 output variable, each containing a similar number of signal decays. This binning scheme enhances the sensitivity while giving stable fit performance.

The B candidate mass shapes in the fit to $B^+ \rightarrow D^- K^+ \pi^+$ candidates are modeled in the same way as described in Ref. [11]. The signal is described by the sum of two Crystal Ball (CB) [26] functions, with a common mean. The tails on both sides of the peak are described by parameters that are fixed to values found in fits to simulated samples. Components are included for combinatorial background, modeled with an exponential shape, and for partially reconstructed backgrounds from $B^+ \rightarrow D^{*-} K^+ \pi^+$ decays and misidentified $B^+ \rightarrow D^{(*)-} \pi^+ \pi^+$ and $B^+ \rightarrow D_s^- K^+ \pi^+$ decays, for which nonparametric descriptions are determined from simulation. Data-driven estimates of the misidentification probabilities, the phase-space distributions of the $B^+ \rightarrow D^{(*)-} \pi^+ \pi^+$ decays [27,28] and the relative branching fractions of the $B^+ \rightarrow D^- \pi^+ \pi^+$ and $B^+ \rightarrow D^{*-} \pi^+ \pi^+$ modes [27,29] are used to obtain these shapes. For signal and partially reconstructed and combinatorial background components, the relative yields

in each NN2 bin are free parameters of the fit, while those of misidentified $B^+ \rightarrow D^{(*)-} \pi^+ \pi^+$ and $B^+ \rightarrow D_s^- K^+ \pi^+$ decays are taken to be the same as for signal decays, since their NN2 responses are expected to be very similar.

A total of 25 parameters are determined from the fit to the $D^- K^+ \pi^+$ sample. These include yields of $B^+ \rightarrow D^- K^+ \pi^+$ decays, the combinatorial background, the partially reconstructed background, and the $B^+ \rightarrow D^{(*)-} \pi^+ \pi^+$ and $B^+ \rightarrow D_s^- K^+ \pi^+$ misidentified backgrounds. For the signal category, and for combinatorial and partially reconstructed backgrounds, the fractional yields f_i of each component in NN2 bins 1–5 are free parameters, with the fraction in bin 6 determined as $f_6 = 1 - \sum_{i=1}^5 f_i$. In addition, the exponential slope parameter of the combinatorial background and parameters of the signal invariant mass shape (the peak position, the width of the core CB function, the relative normalization and ratio of the CB widths) are allowed to vary. Figure 2 shows the combined B^+ candidate mass distribution of all NN2 bins weighted by $S/(S+B)$, where S and B are the fitted signal and background yields within $\pm 2.5\sigma_{\text{CB}}$ of the signal peak position and σ_{CB} is the width of the core CB function. The fit results are summarized in Table I.

The model for the fit to $B^+ \rightarrow D^+ K^+ \pi^-$ candidates is similar to that for the $B^+ \rightarrow D^- K^+ \pi^+$ case. The functional forms for the mass shapes for signal and combinatorial background are identical. The signal peak position, the width of the core CB function, and the fractional yields in each NN2 bin are fixed to the values obtained from the $B^+ \rightarrow D^- K^+ \pi^+$ fit. A component is included for partially reconstructed background, which is likely to be dominated by $\bar{B}_s^0 \rightarrow D^+ K^+ \pi^- \pi^-$ decays; although this channel is unobserved, it is expected to be a sizable source of background based on studies of similar decay modes [30–32]. As the resonant structure of this mode is unknown, its mass shape is modeled using a combination of simulated samples generated with various $D^+ \pi^-$, $D^+ K^+ \pi^-$ and $K^+ \pi^-$ resonances and nonresonant amplitudes. The unknown structure of this background could cause some disagreement between data and the fit result at low $m(D^+ K^+ \pi^-)$. The fractional yields in each NN2 bin are fixed to be the same as those for partially reconstructed backgrounds in the $B^+ \rightarrow D^- K^+ \pi^+$ fit. Potential partially reconstructed background from B^0 and B^+ decays with a missing pion hardly enter the fit region; any residual contributions are absorbed in the $\bar{B}_s^0 \rightarrow D^+ K^+ \pi^- \pi^-$ mass shape.




FIG. 2. Weighted invariant mass distribution of candidates in the (left) $B^+ \rightarrow D^- K^+ \pi^+$ and (right) $B^+ \rightarrow D^+ K^+ \pi^-$ data samples. Data points and fit functions are weighted as described in the text. The components are as detailed in the legend.

There remain 11 parameters that are varied in the fit to the $D^+ K^+ \pi^-$ sample: the yields for $B^+ \rightarrow D^+ K^+ \pi^-$ decays, combinatorial and partially reconstructed backgrounds; the fractional yields of the combinatorial background in each NN2 bin; the exponential slope parameter of the combinatorial background and the relative normalization and ratio of widths of the two CB functions. The results of this fit are summarized in Table I and shown in Fig. 2. The statistical significance of the $D^+ K^+ \pi^-$ peak, obtained from the square root of twice the change in negative log likelihood from the value obtained in a fit with zero signal yield, is 11σ .

Systematic uncertainties on the ratio $\mathcal{B}(B^+ \rightarrow D^+ K^+ \pi^-) / \mathcal{B}(B^+ \rightarrow D^- K^+ \pi^+)$ arise due to approximations made in the signal and background shapes used in the fit and uncertainties in the relative efficiencies. The largest uncertainties are associated with the particle identification and hardware trigger efficiencies (5.9%), the modeling of the combinatorial background in B candidate mass and NN2 bins (4.4%) and the NN2 distributions of signal and partially reconstructed background (4.2%). Other sources, including the modeling of the $B_s^0 \rightarrow D^+ K^- \pi^+ \pi^-$ background and potential biases that are either intrinsic to the fit procedure or related to the treatment of multiple candidates, contribute systematic uncertainties of 2.0% or less.

With all sources combined in quadrature, the total systematic uncertainty on the ratio of branching fractions is found to be 9.0%. The likelihood function is convolved

TABLE I. Yields and statistical uncertainties obtained from fits to the $D^- K^+ \pi^+$ and $D^+ K^+ \pi^-$ data samples.

	$D^- K^+ \pi^+$	$D^+ K^+ \pi^-$
$B^+ \rightarrow D^+ K^+ \pi^-$...	164 ± 21
$B^+ \rightarrow D^- K^+ \pi^+$	3101 ± 66	...
Combinatorial background	3710 ± 110	5945 ± 89
Partially reconstructed background	1676 ± 57	1425 ± 54
$B^+ \rightarrow D^{(*)-} \pi^+ \pi^+$	548 ± 67	...
$B^+ \rightarrow D_s^- K^+ \pi^+$	342 ± 42	...

with a Gaussian of width corresponding to the size of the systematic uncertainties that affect the signal yield, and the total significance of the signal is found to be 8σ .

The relative branching fraction of $B^+ \rightarrow D^+ K^+ \pi^-$ and $B^+ \rightarrow D^- K^+ \pi^+$ decays is determined from

$$\frac{\mathcal{B}(B^+ \rightarrow D^+ K^+ \pi^-)}{\mathcal{B}(B^+ \rightarrow D^- K^+ \pi^+)} = \frac{N^{\text{corr}}(B^+ \rightarrow D^+ K^+ \pi^-)}{N^{\text{corr}}(B^+ \rightarrow D^- K^+ \pi^+)}, \quad (1)$$

where the efficiency-corrected yield is $N^{\text{corr}} = \sum_i W_i / \epsilon_i$. Here the index i runs over all candidates in the fit range, W_i is the signal weight for candidate i , determined using the *sPlot* procedure [33], from the fits shown in Fig. 2, and ϵ_i is the efficiency for candidate i as a function of its Dalitz plot position.

The average efficiencies are defined as $\bar{\epsilon} = N / N^{\text{corr}} = \sum_i W_i / N^{\text{corr}}$ and are found to be $\bar{\epsilon}(B^+ \rightarrow D^+ K^+ \pi^-) = (0.057 \pm 0.014)\%$ and $\bar{\epsilon}(B^+ \rightarrow D^- K^+ \pi^+) = (0.079 \pm 0.003)\%$. These values include contributions from the LHCb detector acceptance, selection and trigger. The trigger efficiency and most selection efficiencies are calculated from simulated samples with data-driven corrections applied, while the particle identification efficiency is measured from a data control sample [34]. The difference between the efficiencies is mainly caused by the different Dalitz plot distributions of the data in each channel.

From Eq. (1), the ratio of branching fractions is determined to be

$$\frac{\mathcal{B}(B^+ \rightarrow D^+ K^+ \pi^-)}{\mathcal{B}(B^+ \rightarrow D^- K^+ \pi^+)} = 0.073 \pm 0.012(\text{stat}) \pm 0.007(\text{syst}).$$

Taking $\mathcal{B}(B^+ \rightarrow D^- K^+ \pi^+) = (7.31 \pm 0.19 \pm 0.22 \pm 0.39) \times 10^{-5}$ [11] gives

$$\mathcal{B}(B^+ \rightarrow D^+ K^+ \pi^-) = (5.31 \pm 0.90 \pm 0.48 \pm 0.35) \times 10^{-6},$$

where the third uncertainty is from $\mathcal{B}(B^+ \rightarrow D^- K^+ \pi^+)$, which arises mainly from the precision with which $\mathcal{B}(B^+ \rightarrow D^- \pi^+ \pi^+)$ [29] is known.




FIG. 3. Background-subtracted Dalitz plot distribution of $B^+ \rightarrow D^+K^+\pi^-$ candidates in the region $5260 < m(D^+K^+\pi^-) < 5310$ MeV/ c^2 from all NN2 bins. Areas of boxes are proportional to signal yields. Negative yields are indicated by crosses.

The Dalitz plot distribution of $B^+ \rightarrow D^+K^+\pi^-$ candidates in the region $5260 < m(D^+K^+\pi^-) < 5310$ MeV/ c^2 is shown in Fig. 3. Combinatorial background has been subtracted using the distribution of candidates in a sideband [$5400 < m(D^+K^+\pi^-) < 5800$ MeV/ c^2], while the signal region has been chosen to minimize the $\bar{B}_s^0 \rightarrow D^+K^+\pi^-\pi^-$ background contribution. The Dalitz plot variables are calculated with a constraint imposed on the B mass; the combinatorial background distribution is not significantly distorted by this procedure. Some excesses are seen at low $m(D\pi)$ and low $m(K\pi)$, but these do not appear to be from narrow structures; rather, there seems to be a broad S -wave $D\pi$ contribution. The apparent structure at high $m(D\pi)$ may arise from imperfect background subtraction.

Although the $B^+ \rightarrow D^+K^+\pi^-$ yield is not sufficient for a Dalitz plot analysis, it is possible to gain information about the contributions from narrow resonances. Two-body mass requirements can reduce the contributions from other intermediate states, but not to a negligible level. Therefore it is necessary to use the angular decay distributions to isolate particular resonances. The $B^+ \rightarrow D^-K^+\pi^+$ normalization mode is again used to reduce potential sources of systematic bias.

Contributions from different partial waves can be determined by weighting the data according to the value of the Legendre polynomial of order L , P_L , evaluated as a function of the cosine of the helicity angle of the $K^+\pi^-$ or $D^+\pi^-$ system. The helicity angle is defined as the angle between the momentum vectors of the pion and the B^+ candidate in the $K^+\pi^-$ or $D^+\pi^-$ rest frame. Event-by-event efficiency corrections, determined as a function of Dalitz plot position, are also applied. The helicity angles and two-body invariant masses are calculated with a constraint on the B mass imposed on the decay chain. If only resonances up to spin J_{\max} are present in a certain mass region, the $P_{2J_{\max}}$ moment will isolate the highest spin state. Thus, in

the limit that only $D^+\pi^-$ resonances contribute, weighting by P_4 will isolate the $D_2^*(2460)^0$ component, as shown in Ref. [11], where a more detailed description of contributions to each moment can be found. Similarly, at low $m(K^+\pi^-)$, weighting by P_2 can be used to determine the contribution from the $K^*(892)^0$ resonance. Higher moments may be present, due to tails of higher spin resonances or reflections from resonances in the other two-particle combination; these will also cause an excess of events in regions away from the resonance peak and therefore can be accounted for by sideband subtraction.

Candidates are selected within regions corresponding to approximately $\pm 2\Gamma$, where Γ is the natural width [29], around the peaks of the $D_2^*(2460)^0$ resonance in $m(D^\pm\pi^\mp)$ and of the $K^*(892)^0$ resonance in $m(K^+\pi^-)$. The data are efficiency corrected and weighted according to the corresponding Legendre polynomial functions. Yields, denoted \tilde{N} , are then obtained from binned minimum χ^2 fits to the B^+ candidate mass distribution. A variable-width binning scheme is used with bin widths chosen to avoid empty bins. The same procedure is applied for candidates in low and high sideband regions, between about 3Γ and 5Γ from the peak. For the normalization of the search for $D^+K^*(892)^0$ decays, the full efficiency-corrected $D^-K^+\pi^+$ sample is used without weighting by angular momentum. The results are used to measure the ratios of branching fractions

$$\frac{\mathcal{B}(B^+ \rightarrow D_2^*(2460)^0 K^+)}{\mathcal{B}(B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+)} \equiv (r_B(D_2^*(2460)K^+))^2 = \frac{\tilde{N}^{\text{corr}}(B^+ \rightarrow D_2^*(2460)^0 K^+)}{\tilde{N}^{\text{corr}}(B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+)}, \quad (2)$$

$$\frac{\mathcal{B}(B^+ \rightarrow D^+K^*(892)^0 \rightarrow D^+K^+\pi^-)}{\mathcal{B}(B^+ \rightarrow D^-K^+\pi^+)} = \frac{\tilde{N}^{\text{corr}}(B^+ \rightarrow D^+K^*(892)^0) \cdot (\frac{5}{2})}{\tilde{N}^{\text{corr}}(B^+ \rightarrow D^-K^+\pi^+) \epsilon(K^*(892)^0)}, \quad (3)$$

where \tilde{N}^{corr} are the yields obtained from the fit after accounting for subtraction of higher moments as estimated from the sideband regions. In Eq. (3) the correction of $\frac{5}{2}$ arises from the normalization of the Legendre polynomial functions and the factor of $\epsilon(K^*(892)^0) = 0.857 \pm 0.006$ is due to the efficiency of the $K^*(892)^0$ signal region [$801.0 < m(K^+\pi^-) < 990.6$ MeV/ c^2] requirement. All efficiency, D_2^* branching fraction and normalization effects cancel in Eq. (2).

The fit models used are based on those described above, but with some important simplifications. The angular weighting by P_2 or P_4 significantly reduces the combinatorial background, and therefore candidates in all NN2 bins are combined; moreover a linear shape is used instead of an




FIG. 4. Results of binned minimum χ^2 fits to the B^+ candidate invariant mass distributions of (left) $B^+ \rightarrow D^- K^+ \pi^+$ and (right) $B^+ \rightarrow D^+ K^+ \pi^-$ candidates with efficiency corrections and angular weights applied. Candidates in the ranges (top) $2217.6 < m(D^\pm \pi^\mp) < 2315.6 \text{ MeV}/c^2$ and (bottom right) $801.0 < m(K^+ \pi^-) < 990.6 \text{ MeV}/c^2$ are shown, while in the bottom left plot the whole $m(K^+ \pi^+)$ range is included. The components are as described in the legend.

exponential function in order to allow for the possibility that the weighted background can fluctuate to negative values. The $B^+ \rightarrow D^{(*)-} \pi^+ \pi^+$ and $\bar{B}_s^0 \rightarrow D^+ K^+ \pi^- \pi^-$ background shapes are given by nonparametric functions obtained from simulated samples with angular moment weighting applied. No component is included for misidentified $B^+ \rightarrow D_s^- K^+ \pi^+$ decays, as it is found to be removed by the weighting procedure.

The analysis method is validated using the $B^+ \rightarrow D^- K^+ \pi^+$ channel and simulated pseudo-experiments. The fit fraction for $B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+$ decays obtained from a full Dalitz plot analysis in Ref. [11] is reproduced within the expected range. The procedure is tested by searching for a fake K^* resonance in $m(K^+ \pi^+)$, and the yield is found to be consistent with zero.

The results of the fits to P_4 -weighted and efficiency-corrected $B^+ \rightarrow D^+ K^+ \pi^-$ and $B^+ \rightarrow D^- K^+ \pi^+$ data samples in the $\bar{D}_2^*(2460)^0$ resonance region are shown in Fig. 4. The procedure isolates the $B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+$ decay, as expected, but no evidence is seen for the suppressed $B^+ \rightarrow D_2^*(2460)^0 K^+$ channel. The corresponding fits for the $B^+ \rightarrow D^+ K^*(892)^0$ search are also shown in Fig. 4; there is no evidence for this decay. The yields are given in Table II.

Systematic uncertainties arise due to the fit models and background subtraction used to determine \tilde{N}^{corr} in Eqs. (2) and (3). The uncertainties are evaluated from the effects on the yields of the following variations: the combinatorial background shape is changed from linear to flat; the

TABLE II. Results of the binned minimum χ^2 fits to efficiency-corrected B^+ candidate invariant mass distributions in each resonance region and with weighting according to angular distributions as described in the text.

	Lower Sideband	Signal Region	Upper Sideband	\tilde{N}^{corr}
$\tilde{N}(B^+ \rightarrow D_2^*(2460)^0 K^+)$	-200 ± 2500	500 ± 3000	200 ± 2200	500 ± 4500
$\tilde{N}(B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+)$	28000 ± 14000	293000 ± 24000	-600 ± 4200	266000 ± 28000
$\tilde{N}(B^+ \rightarrow D^+ K^*(892)^0)$	1700 ± 1900	-3000 ± 5000	9500 ± 4000	-14000 ± 7000
$\tilde{N}(B^+ \rightarrow D^- K^+ \pi^+)$...	4670000 ± 110000

R. AAIJ *et al.*

PHYSICAL REVIEW D **93**, 051101(R) (2016)

$\bar{B}_s^0 \rightarrow D^+ K^+ \pi^- \pi^-$ background component is removed; all other fit components are varied in the same way as described previously. The limited precision of the knowledge of the efficiencies as functions of Dalitz plot position also causes a small uncertainty. An uncertainty is assigned due to the effect of changing the sideband regions from the default of $3\Gamma \leftrightarrow 5\Gamma$ to $4\Gamma \leftrightarrow 6\Gamma$. The uncertainty in $\epsilon(K^*(892)^0)$ of Eq. (3) is also accounted for. The total systematic uncertainty is obtained by combining all sources in quadrature.

The ratio of branching fractions is thus measured to be

$$\frac{\mathcal{B}(B^+ \rightarrow D_2^*(2460)^0 K^+)}{\mathcal{B}(B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+)} = 0.002 \pm 0.015(\text{stat}) \pm 0.005(\text{syst}),$$

which in turn gives

$$r_B(D_2^*(2460)K^+) = 0.04 \pm 0.18(\text{stat}) \pm 0.06(\text{syst}).$$

Assuming Gaussian uncertainties, upper limits at 90(95)% confidence level (CL) are obtained by integrating the likelihood in the region of positive branching fraction,

$$(r_B(D_2^*(2460)K^+))^2 < 0.027(0.033) \quad \text{and} \\ r_B(D_2^*(2460)K^+) < 0.30(0.36).$$

The result for $(r_B(D_2^*(2460)K^+))^2$ and the product branching fraction $\mathcal{B}(B^+ \rightarrow \bar{D}_2^*(2460)^0 K^+) \times \mathcal{B}(\bar{D}_2^*(2460)^0 \rightarrow D^- \pi^+) = (23.2 \pm 1.1 \pm 0.6 \pm 1.0 \pm 1.6) \times 10^{-4}$ at 90(95)% C.L. [11] give

$$\mathcal{B}(B^+ \rightarrow D_2^*(2460)^0 K^+) \times \mathcal{B}(D_2^*(2460)^0 \rightarrow D^+ \pi^-) \\ = (0.4 \pm 3.5 \pm 1.1 \pm 0.1) \times 10^{-5}, \\ < 6.3(7.5) \times 10^{-5} \quad \text{at 90(95)\% CL.}$$

These are the first experimental results on this decay mode. Similarly for $B^+ \rightarrow D^+ K^*(892)^0 \rightarrow D^+ K^+ \pi^-$,

$$\frac{\mathcal{B}(B^+ \rightarrow D^+ K^*(892)^0 \rightarrow D^+ K^+ \pi^-)}{\mathcal{B}(B^+ \rightarrow D^- K^+ \pi^+)} \\ = -0.0079 \pm 0.0039(\text{stat}) \pm 0.0028(\text{syst}), \\ < 0.0044(0.0055) \quad \text{at 90(95)\% C.L.}$$

The measured value $\mathcal{B}(B^+ \rightarrow D^- K^+ \pi^+) = (7.31 \pm 0.19 \pm 0.22 \pm 0.39) \times 10^{-5}$ [11] and the isospin relation $\mathcal{B}(K^*(892)^0 \rightarrow K^+ \pi^-) = \frac{2}{3}$ give

$$\mathcal{B}(B^+ \rightarrow D^+ K^*(892)^0) = (-8.7 \pm 4.3 \pm 3.1 \pm 0.4) \times 10^{-7}, \\ < 4.9(6.1) \times 10^{-7} \quad \text{at 90(95)\% C.L.,}$$

where the third uncertainty is due to the normalization channel branching fraction. This result supersedes the previous limit, which was obtained with a subset of the data [12].

In summary, the rare $B^+ \rightarrow D^+ K^+ \pi^-$ decay has been observed for the first time with 8σ significance, based on a data sample of 3.0 fb^{-1} of pp collision data collected with the LHCb detector. The Dalitz plot appears to be dominated by broad structures. Searches for $B^+ \rightarrow D_2^*(2460)^0 K^+$ and $B^+ \rightarrow D^+ K^*(892)^0$ decays have been carried out by weighting the data according to the decay angle distributions, but no significant signals are seen. These results indicate that further studies, with larger data samples, of the Dalitz plot distribution of this mode will be of interest to understand the potential for a measurement of γ from $B^+ \rightarrow DK^+ \pi^0$ decays.

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); FOM and NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinCo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (U.S.). We are indebted to the communities behind the multiple open source software packages on which we depend. We are also thankful for the computing resources and the access to software R&D tools provided by Yandex LLC (Russia). Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR (Russia), GVA, XuntaGal and GENCAT (Spain), The Royal Society and Royal Commission for the Exhibition of 1851 (United Kingdom).

- [1] N. Cabibbo, Unitary Symmetry and Leptonic Decays, *Phys. Rev. Lett.* **10**, 531 (1963).
- [2] M. Kobayashi and T. Maskawa, CP violation in the renormalizable theory of weak interaction, *Prog. Theor. Phys.* **49**, 652 (1973).
- [3] M. Bona *et al.* (UTfit Collaboration), The 2004 UTfit Collaboration report on the status of the unitarity triangle in the standard model, *J. High Energy Phys.* **07** (2005) 028.
- [4] J. Charles, A. Höcker, H. Lacker, S. Laplace, F. R. Diberder, J. Malclés, J. Ocariz, M. Pivk, and L. Roos (CKMfitter Group), CP violation and the CKM matrix: Assessing the impact of the asymmetric B factories, *Eur. Phys. J. C* **41**, 1 (2005).
- [5] R. Aaij (LHCb Collaboration), Improved constraints on γ : CKM2014 update, Report No. LHCb-CONF-2014-004.
- [6] M. Gronau and D. London, How to determine all the angles of the unitarity triangle from $B^0 \rightarrow DK_S^0$ and $B_s^0 \rightarrow D\phi$, *Phys. Lett. B* **253**, 483 (1991).
- [7] M. Gronau and D. Wyler, On determining a weak phase from charged B decay asymmetries, *Phys. Lett. B* **265**, 172 (1991).
- [8] D. Atwood, I. Dunietz, and A. Soni, Enhanced CP Violation with $B \rightarrow KD^0(\bar{D}^0)$ Modes and Extraction of the CKM Angle γ , *Phys. Rev. Lett.* **78**, 3257 (1997).
- [9] D. Atwood, I. Dunietz, and A. Soni, Improved methods for observing CP violation in $B^\pm \rightarrow K^\pm D$ and measuring the CKM phase γ , *Phys. Rev. D* **63**, 036005 (2001).
- [10] N. Sinha, Determining γ using $B \rightarrow D^{**}K$, *Phys. Rev. D* **70**, 097501 (2004).
- [11] R. Aaij *et al.* (LHCb Collaboration), First observation and amplitude analysis of the $B^- \rightarrow D^+ K^- \pi^-$ decay, *Phys. Rev. D* **91**, 092002 (2015).
- [12] R. Aaij *et al.* (LHCb Collaboration), First evidence for the annihilation decay mode $B^+ \rightarrow D_s^+ \phi$, *J. High Energy Phys.* **02** (2013) 043.
- [13] A. A. Alves, Jr. *et al.* (LHCb Collaboration), The LHCb detector at the LHC, *J. Instrum.* **3**, S08005 (2008).
- [14] R. Aaij *et al.* (LHCb Collaboration), LHCb detector performance, *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
- [15] T. Sjöstrand, S. Mrenna, and P. Skands, A brief introduction to PYTHIA 8.1, *Comput. Phys. Commun.* **178**, 852 (2008); PYTHIA 6.4 physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [16] I. Belyaev *et al.*, Handling of the generation of primary events in Gauss, the LHCb simulation framework, *J. Phys. Conf. Ser.* **331**, 032047 (2011).
- [17] D. J. Lange, The EvtGen particle decay simulation package, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [18] S. Agostinelli *et al.* (Geant4 Collaboration), Geant4: A simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003); J. Allison *et al.* (Geant4 Collaboration), Geant4 developments and applications, *IEEE Trans. Nucl. Sci.* **53**, 270 (2006).
- [19] P. Golonka and Z. Was, PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays, *Eur. Phys. J. C* **45**, 97 (2006).
- [20] M. Clemencic, G. Corti, S. Easo, C. R. Jones, S. Miglioranza, M. Pappagallo, and P. Robbe, The LHCb simulation application, Gauss: Design, evolution and experience, *J. Phys. Conf. Ser.* **331**, 032023 (2011).
- [21] M. Feindt and U. Kerzel, The NeuroBayes neural network package, *Nucl. Instrum. Methods Phys. Res., Sect. A* **559**, 190 (2006).
- [22] R. Aaij *et al.* (LHCb Collaboration), Measurement of the $B_s^0 \rightarrow \mu^+ \mu^-$ Branching Fraction and Search for $B^0 \rightarrow \mu^+ \mu^-$ Decays at the LHCb Experiment, *Phys. Rev. Lett.* **111**, 101805 (2013).
- [23] R. Aaij *et al.* (LHCb Collaboration), Search for the lepton flavour violating decay $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, *J. High Energy Phys.* **02** (2015) 121.
- [24] R. Aaij *et al.* (LHCb Collaboration), Search for the decay $B_s^0 \rightarrow \bar{D}^0 f_0(980)$, *J. High Energy Phys.* **08** (2015) 005.
- [25] R. Aaij *et al.* (LHCb Collaboration), Search for the lepton-flavour violating decay $D^0 \rightarrow e^\pm \mu^\mp$, *Phys. Lett. B* **754**, 167 (2016).
- [26] T. Skwarnicki, Ph. D. thesis, Institute of Nuclear Physics, Krakow, [Report No. DESY-F31-86-02, 1986].
- [27] K. Abe *et al.* (Belle Collaboration), Study of $B^- \rightarrow D^{*0} \pi^- (D^{*0} \rightarrow D^{(*)+} \pi^-)$ decays, *Phys. Rev. D* **69**, 112002 (2004).
- [28] B. Aubert *et al.* (BABAR Collaboration), Dalitz plot analysis of $B^- \rightarrow D^+ \pi^- \pi^-$, *Phys. Rev. D* **79**, 112004 (2009).
- [29] K. A. Olive *et al.* (Particle Data Group), Review of particle physics, *Chin. Phys. C* **38**, 090001 (2014).
- [30] R. Aaij *et al.* (LHCb Collaboration), First observation of the decays $\bar{B}_{(s)}^0 \rightarrow D_s^+ K^- \pi^+ \pi^-$ and $\bar{B}_s^0 \rightarrow D_{s1}(2536)^+ \pi^-$, *Phys. Rev. D* **86**, 112005 (2012).
- [31] R. Aaij *et al.* (LHCb Collaboration), Observation of Overlapping Spin-1 and Spin-3 $\bar{D}^0 K^-$ Resonances at Mass $2.86 \text{ GeV}/c^2$, *Phys. Rev. Lett.* **113**, 162001 (2014).
- [32] R. Aaij *et al.* (LHCb Collaboration), Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays, *Phys. Rev. D* **90**, 072003 (2014).
- [33] M. Pivk and F. R. Le Diberder, sPlot: A statistical tool to unfold data distributions, *Nucl. Instrum. Methods Phys. Res., Sect. A* **555**, 356 (2005).
- [34] M. Adinolfi *et al.*, Performance of the LHCb RICH detector at the LHC, *Eur. Phys. J. C* **73**, 2431 (2013).

R. Aaij,³⁹ C. Abellán Beteta,⁴¹ B. Adeva,³⁸ M. Adinolfi,⁴⁷ A. Affolder,⁵³ Z. Ajaltouni,⁵ S. Akar,⁶ J. Albrecht,¹⁰ F. Alessio,³⁹ M. Alexander,⁵² S. Ali,⁴² G. Alkhazov,³¹ P. Alvarez Cartelle,⁵⁴ A. A. Alves Jr.,⁵⁸ S. Amato,² S. Amerio,²³ Y. Amhis,⁷ L. An,^{3,40} L. Anderlini,¹⁸ G. Andreassi,⁴⁰ M. Andreotti,^{17,g} J. E. Andrews,⁵⁹ R. B. Appleby,⁵⁵ O. Aquines Gutierrez,¹¹ F. Archilli,³⁹ P. d'Argent,¹² A. Artamonov,³⁶ M. Artuso,⁶⁰ E. Aslanides,⁶ G. Auriemma,^{26,n} M. Baalouch,⁵ S. Bachmann,¹² J. J. Back,⁴⁹ A. Badalov,³⁷ C. Baesso,⁶¹ W. Baldini,^{17,39} R. J. Barlow,⁵⁵ C. Barschel,³⁹

S. Barsuk,⁷ W. Barter,³⁹ V. Batozskaya,²⁹ V. Battista,⁴⁰ A. Bay,⁴⁰ L. Beaucourt,⁴ J. Beddow,⁵² F. Bedeschi,²⁴ I. Bediaga,¹ L. J. Bel,⁴² V. Bellee,⁴⁰ N. Belloli,^{21,k} I. Belyaev,³² E. Ben-Haim,⁸ G. Bencivenni,¹⁹ S. Benson,³⁹ J. Benton,⁴⁷ A. Berezhnoy,³³ R. Bernet,⁴¹ A. Bertolin,²³ M. -O. Bettler,³⁹ M. van Beuzekom,⁴² S. Bifani,⁴⁶ P. Billoir,⁸ T. Bird,⁵⁵ A. Birnkraut,¹⁰ A. Bizzeti,^{18,i} T. Blake,⁴⁹ F. Blanc,⁴⁰ J. Blouw,¹¹ S. Blusk,⁶⁰ V. Bocci,²⁶ A. Bondar,³⁵ N. Bondar,^{31,39} W. Bonivento,¹⁶ S. Borghi,⁵⁵ M. Borisyak,⁶⁶ M. Borsato,³⁸ T. J. V. Bowcock,⁵³ E. Bowen,⁴¹ C. Bozzi,^{17,39} S. Braun,¹² M. Britsch,¹² T. Britton,⁶⁰ J. Brodzicka,⁵⁵ N. H. Brook,⁴⁷ E. Buchanan,⁴⁷ C. Burr,⁵⁵ A. Bursche,⁴¹ J. Buytaert,³⁹ S. Cadeddu,¹⁶ R. Calabrese,^{17,g} M. Calvi,^{21,k} M. Calvo Gomez,^{37,p} P. Campana,¹⁹ D. Campora Perez,³⁹ L. Capriotti,⁵⁵ A. Carbone,^{15,e} G. Carboni,^{25,l} R. Cardinale,^{20,j} A. Cardini,¹⁶ P. Carniti,^{21,k} L. Carson,⁵¹ K. Carvalho Akiba,² G. Casse,⁵³ L. Cassina,^{21,k} L. Castillo Garcia,⁴⁰ M. Cattaneo,³⁹ Ch. Cauet,¹⁰ G. Cavallero,²⁰ R. Cenci,^{24,t} M. Charles,⁸ Ph. Charpentier,³⁹ M. Chefdeville,⁴ S. Chen,⁵⁵ S.-F. Cheung,⁵⁶ N. Chiapolini,⁴¹ M. Chrzasczcz,^{41,27} X. Cid Vidal,³⁹ G. Ciezarek,⁴² P. E. L. Clarke,⁵¹ M. Clemencic,³⁹ H. V. Cliff,⁴⁸ J. Closier,³⁹ V. Coco,³⁹ J. Cogan,⁶ E. Cogneras,⁵ V. Cogoni,^{16,f} L. Cojocariu,³⁰ G. Collazuol,^{23,r} P. Collins,³⁹ A. Comerma-Montells,¹² A. Contu,³⁹ A. Cook,⁴⁷ M. Coombes,⁴⁷ S. Coquereau,⁸ G. Corti,³⁹ M. Corvo,^{17,g} B. Couturier,³⁹ G. A. Cowan,⁵¹ D. C. Craik,⁵¹ A. Crocombe,⁴⁹ M. Cruz Torres,⁶¹ S. Cunliffe,⁵⁴ R. Currie,⁵⁴ C. D'Ambrosio,³⁹ E. Dall'Occo,⁴² J. Dalseno,⁴⁷ P. N. Y. David,⁴² A. Davis,⁵⁸ O. De Aguiar Francisco,² K. De Bruyn,⁶ S. De Capua,⁵⁵ M. De Cian,¹² J. M. De Miranda,¹ L. De Paula,² P. De Simone,¹⁹ C. -T. Dean,⁵² D. Decamp,⁴ M. Deckenhoff,¹⁰ L. Del Buono,⁸ N. Déléage,⁴ M. Demmer,¹⁰ D. Derkach,⁶⁶ O. Deschamps,⁵ F. Dettori,³⁹ B. Dey,²² A. Di Canto,³⁹ F. Di Ruscio,²⁵ H. Dijkstra,³⁹ S. Donleavy,⁵³ F. Dordei,³⁹ M. Dorigo,⁴⁰ A. Dosil Suárez,³⁸ A. Dovbnya,⁴⁴ K. Dreimanis,⁵³ L. Dufour,⁴² G. Dujany,⁵⁵ K. Dungs,³⁹ P. Durante,³⁹ R. Dzhelyadin,³⁶ A. Dziurda,²⁷ A. Dzyuba,³¹ S. Easo,^{50,39} U. Egede,⁵⁴ V. Egorychev,³² S. Eidelman,³⁵ S. Eisenhardt,⁵¹ U. Eitschberger,¹⁰ R. Ekelhof,¹⁰ L. Eklund,⁵² I. El Rifai,⁵ Ch. Elsasser,⁴¹ S. Ely,⁶⁰ S. Esen,¹² H. M. Evans,⁴⁸ T. Evans,⁵⁶ M. Fabianska,²⁷ A. Falabella,¹⁵ C. Färber,³⁹ N. Farley,⁴⁶ S. Farry,⁵³ R. Fay,⁵³ D. Ferguson,⁵¹ V. Fernandez Albor,³⁸ F. Ferrari,¹⁵ F. Ferreira Rodrigues,¹ M. Ferro-Luzzi,³⁹ S. Filippov,³⁴ M. Fiore,^{17,39,g} M. Fiorini,^{17,g} M. Firlej,²⁸ C. Fitzpatrick,⁴⁰ T. Fiutowski,²⁸ F. Fleuret,^{7,b} K. Fohl,³⁹ P. Fol,⁵⁴ M. Fontana,¹⁶ F. Fontanelli,^{20,j} D. C. Forshaw,⁶⁰ R. Forty,³⁹ M. Frank,³⁹ C. Frei,³⁹ M. Frosini,¹⁸ J. Fu,²² E. Furfaro,^{25,l} A. Gallas Torreira,³⁸ D. Galli,^{15,e} S. Gallorini,²³ S. Gambetta,⁵¹ M. Gandelman,² P. Gandini,⁵⁶ Y. Gao,³ J. García Pardiñas,³⁸ J. Garra Tico,⁴⁸ L. Garrido,³⁷ D. Gascon,³⁷ C. Gaspar,³⁹ R. Gauld,⁵⁶ L. Gavardi,¹⁰ G. Gazzoni,⁵ D. Gerick,¹² E. Gersabeck,¹² M. Gersabeck,⁵⁵ T. Gershon,⁴⁹ Ph. Ghez,⁴ S. Gianì,⁴⁰ V. Gibson,⁴⁸ O. G. Girard,⁴⁰ L. Giubega,³⁰ V. V. Gligorov,³⁹ C. Göbel,⁶¹ D. Golubkov,³² A. Golutvin,^{54,39} A. Gomes,^{1,a} C. Gotti,^{21,k} M. Grabalosa Gándara,⁵ R. Graciani Diaz,³⁷ L. A. Granado Cardoso,³⁹ E. Graugés,³⁷ E. Graverini,⁴¹ G. Graziani,¹⁸ A. Greco,³⁰ E. Greening,⁵⁶ P. Griffith,⁴⁶ L. Grillo,¹² O. Grünberg,⁶⁴ B. Gui,⁶⁰ E. Gushchin,³⁴ Yu. Guz,^{36,39} T. Gys,³⁹ T. Hadavizadeh,⁵⁶ C. Hadjivasiliou,⁶⁰ G. Haefeli,⁴⁰ C. Haen,³⁹ S. C. Haines,⁴⁸ S. Hall,⁵⁴ B. Hamilton,⁵⁹ X. Han,¹² S. Hansmann-Menzemer,¹² N. Harnew,⁵⁶ S. T. Harnew,⁴⁷ J. Harrison,⁵⁵ J. He,³⁹ T. Head,⁴⁰ V. Heijne,⁴² A. Heister,⁹ K. Hennessy,⁵³ P. Henrard,⁵ L. Henry,⁸ J. A. Hernando Morata,³⁸ E. van Herwijnen,³⁹ M. Heß,⁶⁴ A. Hicheur,² D. Hill,⁵⁶ M. Hoballah,⁵ C. Hombach,⁵⁵ W. Hulsbergen,⁴² T. Humair,⁵⁴ M. Hushchyn,⁶⁶ N. Hussain,⁵⁶ D. Hutchcroft,⁵³ D. Hynds,⁵² M. Idzik,²⁸ P. Ilten,⁵⁷ R. Jacobsson,³⁹ A. Jaeger,¹² J. Jalocha,⁵⁶ E. Jans,⁴² A. Jawahery,⁵⁹ M. John,⁵⁶ D. Johnson,³⁹ C. R. Jones,⁴⁸ C. Joram,³⁹ B. Jost,³⁹ N. Jurik,⁶⁰ S. Kandybei,⁴⁴ W. Kanso,⁶ M. Karacson,³⁹ T. M. Karbach,^{39,†} S. Karodia,⁵² M. Kecke,¹² M. Kelsey,⁶⁰ I. R. Kenyon,⁴⁶ M. Kenzie,³⁹ T. Ketel,⁴³ E. Khairullin,⁶⁶ B. Khanji,^{21,39,k} C. Khurewathanakul,⁴⁰ T. Kim,⁹ S. Klaver,⁵⁵ K. Klimaszewski,²⁹ O. Kochebina,⁷ M. Kolpin,¹² I. Komarov,⁴⁰ R. F. Koopman,⁴³ P. Koppenburg,^{42,39} M. Kozeiha,⁵ L. Kravchuk,³⁴ K. Kreplin,¹² M. Kreps,⁴⁹ P. Krokovny,³⁵ F. Kruse,¹⁰ W. Krzemien,²⁹ W. Kucewicz,^{27,o} M. Kucharczyk,²⁷ V. Kudryavtsev,³⁵ A. K. Kuonen,⁴⁰ K. Kurek,²⁹ T. Kvaratskheliya,³² D. Lacarrere,³⁹ G. Lafferty,^{55,39} A. Lai,¹⁶ D. Lambert,⁵¹ G. Lanfranchi,¹⁹ C. Langenbruch,⁴⁹ B. Langhans,³⁹ T. Latham,⁴⁹ C. Lazzeroni,⁴⁶ R. Le Gac,⁶ J. van Leerdam,⁴² J. -P. Lees,⁴ R. Lefèvre,⁵ A. Leflat,^{33,39} J. Lefrançois,⁷ E. Lemos Cid,³⁸ O. Leroy,⁶ T. Lesiak,²⁷ B. Leverington,¹² Y. Li,⁷ T. Likhomanenko,^{66,65} M. Liles,⁵³ R. Lindner,³⁹ C. Linn,³⁹ F. Lionetto,⁴¹ B. Liu,¹⁶ X. Liu,³ D. Loh,⁴⁹ I. Longstaff,⁵² J. H. Lopes,² D. Lucchesi,^{23,r} M. Lucio Martinez,³⁸ H. Luo,⁵¹ A. Lupato,²³ E. Luppi,^{17,g} O. Lupton,⁵⁶ A. Lusiani,²⁴ F. Machefert,⁷ F. Maciuc,³⁰ O. Maev,³¹ K. Maguire,⁵⁵ S. Malde,⁵⁶ A. Malinin,⁶⁵ G. Manca,⁷ G. Mancinelli,⁶ P. Manning,⁶⁰ A. Mapelli,³⁹ J. Maratas,⁵ J. F. Marchand,⁴ U. Marconi,¹⁵ C. Marin Benito,³⁷ P. Marino,^{24,39,i} J. Marks,¹² G. Martellotti,²⁶ M. Martin,⁶ M. Martinelli,⁴⁰ D. Martinez Santos,³⁸ F. Martinez Vidal,⁶⁷ D. Martins Tostes,² L. M. Massacrier,⁷ A. Massafferri,¹ R. Matev,³⁹ A. Mathad,⁴⁹ Z. Mathe,³⁹ C. Matteuzzi,²¹ A. Mauri,⁴¹ B. Maurin,⁴⁰ A. Mazurov,⁴⁶ M. McCann,⁵⁴ J. McCarthy,⁴⁶ A. McNab,⁵⁵ R. McNulty,¹³ B. Meadows,⁵⁸ F. Meier,¹⁰ M. Meissner,¹² D. Melnychuk,²⁹ M. Merk,⁴² E. Michielin,²³ D. A. Milanese,⁶³ M. -N. Minard,⁴ D. S. Mitzel,¹²

J. Molina Rodriguez,⁶¹ I. A. Monroy,⁶³ S. Monteil,⁵ M. Morandin,²³ P. Morawski,²⁸ A. Mordà,⁶ M. J. Morello,^{24,t} J. Moron,²⁸ A. B. Morris,⁵¹ R. Mountain,⁶⁰ F. Muheim,⁵¹ D. Müller,⁵⁵ J. Müller,¹⁰ K. Müller,⁴¹ V. Müller,¹⁰ M. Mussini,¹⁵ B. Muster,⁴⁰ P. Naik,⁴⁷ T. Nakada,⁴⁰ R. Nandakumar,⁵⁰ A. Nandi,⁵⁶ I. Nasteva,² M. Needham,⁵¹ N. Neri,²² S. Neubert,¹² N. Neufeld,³⁹ M. Neuner,¹² A. D. Nguyen,⁴⁰ T. D. Nguyen,⁴⁰ C. Nguyen-Mau,^{40,q} V. Niess,⁵ R. Niet,¹⁰ N. Nikitin,³³ T. Nikodem,¹² A. Novoselov,³⁶ D. P. O'Hanlon,⁴⁹ A. Oblakowska-Mucha,²⁸ V. Obraztsov,³⁶ S. Ogilvy,⁵² O. Okhrimenko,⁴⁵ R. Oldeman,^{16,f} C. J. G. Onderwater,⁶⁸ B. Osorio Rodrigues,¹ J. M. Otalora Goicochea,² A. Otto,³⁹ P. Owen,⁵⁴ A. Oyanguren,⁶⁷ A. Palano,^{14,d} F. Palombo,^{22,u} M. Palutan,¹⁹ J. Panman,³⁹ A. Papanestis,⁵⁰ M. Pappagallo,⁵² L. L. Pappalardo,^{17,g} C. Pappenheimer,⁵⁸ W. Parker,⁵⁹ C. Parkes,⁵⁵ G. Passaleva,¹⁸ G. D. Patel,⁵³ M. Patel,⁵⁴ C. Patrignani,^{20,j} A. Pearce,^{55,50} A. Pellegrino,⁴² G. Penso,^{26,m} M. Pepe Altarelli,³⁹ S. Perazzini,^{15,e} P. Perret,⁵ L. Pescatore,⁴⁶ K. Petridis,⁴⁷ A. Petrolini,^{20,j} M. Petruzzo,²² E. Picatoste Olloqui,³⁷ B. Pietrzyk,⁴ M. Pikies,²⁷ D. Pinci,²⁶ A. Pistone,²⁰ A. Piucci,¹² S. Playfer,⁵¹ M. Plo Casasus,³⁸ T. Poikela,³⁹ F. Polci,⁸ A. Poluektov,^{49,35} I. Polyakov,³² E. Polycarpo,² A. Popov,³⁶ D. Popov,^{11,39} B. Popovici,³⁰ C. Potterat,² E. Price,⁴⁷ J. D. Price,⁵³ J. Prisciandaro,³⁸ A. Pritchard,⁵³ C. Prouve,⁴⁷ V. Pugatch,⁴⁵ A. Puig Navarro,⁴⁰ G. Punzi,^{24,s} W. Qian,⁴ R. Quagliani,^{7,47} B. Rachwal,²⁷ J. H. Rademacker,⁴⁷ M. Rama,²⁴ M. Ramos Pernas,³⁸ M. S. Rangel,² I. Raniuk,⁴⁴ N. Rauschmayr,³⁹ G. Raven,⁴³ F. Redi,⁵⁴ S. Reichert,⁵⁵ A. C. dos Reis,¹ V. Renaudin,⁷ S. Ricciardi,⁵⁰ S. Richards,⁴⁷ M. Rihl,³⁹ K. Rinnert,^{53,39} V. Rives Molina,³⁷ P. Robbe,^{7,39} A. B. Rodrigues,¹ E. Rodrigues,⁵⁵ J. A. Rodriguez Lopez,⁶³ P. Rodriguez Perez,⁵⁵ S. Roiser,³⁹ V. Romanovsky,³⁶ A. Romero Vidal,³⁸ J. W. Ronayne,¹³ M. Rotondo,²³ T. Ruf,³⁹ P. Ruiz Valls,⁶⁷ J. J. Saborido Silva,³⁸ N. Sagidova,³¹ B. Saitta,^{16,f} V. Salustino Guimaraes,² C. Sanchez Mayordomo,⁶⁷ B. Sanmartin Sedes,³⁸ R. Santacesaria,²⁶ C. Santamarina Rios,³⁸ M. Santimaria,¹⁹ E. Santovetti,^{25,l} A. Sarti,^{19,m} C. Satriano,^{26,n} A. Satta,²⁵ D. M. Saunders,⁴⁷ D. Savrina,^{32,33} S. Schael,⁹ M. Schiller,³⁹ H. Schindler,³⁹ M. Schlupp,¹⁰ M. Schmelling,¹¹ T. Schmelzer,¹⁰ B. Schmidt,³⁹ O. Schneider,⁴⁰ A. Schopper,³⁹ M. Schubiger,⁴⁰ M. -H. Schune,⁷ R. Schwemmer,³⁹ B. Sciascia,¹⁹ A. Sciubba,^{26,m} A. Semennikov,³² A. Sergi,⁴⁶ N. Serra,⁴¹ J. Serrano,⁶ L. Sestini,²³ P. Seyfert,²¹ M. Shapkin,³⁶ I. Shapoval,^{17,44,g} Y. Shcheglov,³¹ T. Shears,⁵³ L. Shekhtman,³⁵ V. Shevchenko,⁶⁵ A. Shires,¹⁰ B. G. Siddi,¹⁷ R. Silva Coutinho,⁴¹ L. Silva de Oliveira,² G. Simi,^{23,s} M. Sirendi,⁴⁸ N. Skidmore,⁴⁷ T. Skwarnicki,⁶⁰ E. Smith,^{56,50} E. Smith,⁵⁴ I. T. Smith,⁵¹ J. Smith,⁴⁸ M. Smith,⁵⁵ H. Snoek,⁴² M. D. Sokoloff,^{58,39} F. J. P. Soler,⁵² F. Soomro,⁴⁰ D. Souza,⁴⁷ B. Souza De Paula,² B. Spaan,¹⁰ P. Spradlin,⁵² S. Sridharan,³⁹ F. Stagni,³⁹ M. Stahl,³⁹ S. Stahl,³⁹ S. Stefkova,⁵⁴ O. Steinkamp,⁴¹ O. Stenyakin,³⁶ S. Stevenson,⁵⁶ S. Stoica,³⁰ S. Stone,⁶⁰ B. Storaci,⁴¹ S. Stracka,^{24,t} M. Straticiu,³⁰ U. Straumann,⁴¹ L. Sun,⁵⁸ W. Sutcliffe,⁵⁴ K. Swientek,²⁸ S. Swientek,¹⁰ V. Syropoulos,⁴³ M. Szczekowski,²⁹ T. Szumlak,²⁸ S. T'Jampens,⁴ A. Tayduganov,⁶ T. Tekampe,¹⁰ G. Tellarini,^{17,g} F. Teubert,³⁹ C. Thomas,⁵⁶ E. Thomas,³⁹ J. van Tilburg,⁴² V. Tisserand,⁴ M. Tobin,⁴⁰ J. Todd,⁵⁸ S. Tolck,⁴³ L. Tomassetti,^{17,g} D. Tonelli,³⁹ S. Topp-Joergensen,⁵⁶ N. Torr,⁵⁶ E. Tournefier,⁴ S. Tourneur,⁴⁰ K. Trabelsi,⁴⁰ M. Traill,⁵² M. T. Tran,⁴⁰ M. Tresch,⁴¹ A. Trisovic,³⁹ A. Tsaregorodtsev,⁶ P. Tsopelas,⁴² N. Tuning,^{42,39} A. Ukleja,²⁹ A. Ustyuzhanin,^{66,65} U. Uwer,¹² C. Vacca,^{16,39,f} V. Vagnoni,¹⁵ G. Valenti,¹⁵ A. Vallier,⁷ R. Vazquez Gomez,¹⁹ P. Vazquez Regueiro,³⁸ C. Vázquez Sierra,³⁸ S. Vecchi,¹⁷ M. van Veghel,⁴³ J. J. Velthuis,⁴⁷ M. Veltri,^{18,h} G. Veneziano,⁴⁰ M. Vesterinen,¹² B. Viaud,⁷ D. Vieira,² M. Vieites Diaz,³⁸ X. Vilasis-Cardona,^{37,p} V. Volkov,³³ A. Vollhardt,⁴¹ D. Voong,⁴⁷ A. Vorobyev,³¹ V. Vorobyev,³⁵ C. Voß,⁶⁴ J. A. de Vries,⁴² R. Waldi,⁶⁴ C. Wallace,⁴⁹ R. Wallace,¹³ J. Walsh,²⁴ J. Wang,⁶⁰ D. R. Ward,⁴⁸ N. K. Watson,⁴⁶ D. Websdale,⁵⁴ A. Weiden,⁴¹ M. Whitehead,³⁹ J. Wicht,⁴⁹ G. Wilkinson,^{56,39} M. Wilkinson,⁶⁰ M. Williams,³⁹ M. P. Williams,⁴⁶ M. Williams,⁵⁷ T. Williams,⁴⁶ F. F. Wilson,⁵⁰ J. Wimberley,⁵⁹ J. Wishahi,¹⁰ W. Wislicki,²⁹ M. Witek,²⁷ G. Wormser,⁷ S. A. Wotton,⁴⁸ K. Wraight,⁵² S. Wright,⁴⁸ K. Wyllie,³⁹ Y. Xie,⁶² Z. Xu,⁴⁰ Z. Yang,³ J. Yu,⁶² X. Yuan,³⁵ O. Yushchenko,³⁶ M. Zangoli,¹⁵ M. Zavertyaev,^{11,c} L. Zhang,³ Y. Zhang,³ A. Zhelezov,¹² A. Zhokhov,³² L. Zhong,³ V. Zhukov,⁹ and S. Zucchelli¹⁵

(LHCb Collaboration)

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil³Center for High Energy Physics, Tsinghua University, Beijing, China⁴LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

- ⁹*I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ¹⁰*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ¹¹*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ¹²*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ¹³*School of Physics, University College Dublin, Dublin, Ireland*
- ¹⁴*Sezione INFN di Bari, Bari, Italy*
- ¹⁵*Sezione INFN di Bologna, Bologna, Italy*
- ¹⁶*Sezione INFN di Cagliari, Cagliari, Italy*
- ¹⁷*Sezione INFN di Ferrara, Ferrara, Italy*
- ¹⁸*Sezione INFN di Firenze, Firenze, Italy*
- ¹⁹*Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy*
- ²⁰*Sezione INFN di Genova, Genova, Italy*
- ²¹*Sezione INFN di Milano Bicocca, Milano, Italy*
- ²²*Sezione INFN di Milano, Milano, Italy*
- ²³*Sezione INFN di Padova, Padova, Italy*
- ²⁴*Sezione INFN di Pisa, Pisa, Italy*
- ²⁵*Sezione INFN di Roma Tor Vergata, Roma, Italy*
- ²⁶*Sezione INFN di Roma La Sapienza, Roma, Italy*
- ²⁷*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ²⁸*AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ²⁹*National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³⁰*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³¹*Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- ³²*Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- ³³*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- ³⁴*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- ³⁵*Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- ³⁶*Institute for High Energy Physics (IHEP), Protvino, Russia*
- ³⁷*Universitat de Barcelona, Barcelona, Spain*
- ³⁸*Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- ³⁹*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴⁰*Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴¹*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴²*Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ⁴³*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ⁴⁴*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁵*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁶*University of Birmingham, Birmingham, United Kingdom*
- ⁴⁷*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁸*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁴⁹*Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁰*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵¹*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵²*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵³*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁴*Imperial College London, London, United Kingdom*
- ⁵⁵*School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁶*Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁷*Massachusetts Institute of Technology, Cambridge, Massachusetts, USA*
- ⁵⁸*University of Cincinnati, Cincinnati, Ohio, USA*
- ⁵⁹*University of Maryland, College Park, Maryland, USA*
- ⁶⁰*Syracuse University, Syracuse, New York, USA*
- ⁶¹*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil
(associated with Universidade Federal do Rio de Janeiro (UFRJ),
Rio de Janeiro, Brazil)*
- ⁶²*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China
(associated with Center for High Energy Physics, Tsinghua University, Beijing, China)*

⁶³*Departamento de Fisica, Universidad Nacional de Colombia, Bogota, Colombia
(associated with LPNHE, Université Pierre et Marie Curie,
Université Paris Diderot, CNRS/IN2P3, Paris, France)*

⁶⁴*Institut für Physik, Universität Rostock, Rostock, Germany
(associated with Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)*

⁶⁵*National Research Centre Kurchatov Institute, Moscow, Russia
(associated with Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)*

⁶⁶*Yandex School of Data Analysis, Moscow, Russia
(associated with Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)*

⁶⁷*Instituto de Fisica Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain
(associated with Universitat de Barcelona, Barcelona, Spain)*

⁶⁸*Van Swinderen Institute, University of Groningen, Groningen, The Netherlands
(associated with Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands)*

[†]Deceased.

^aAlso at Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil.

^bAlso at Laboratoire Leprince-Ringuet, Palaiseau, France.

^cAlso at P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.

^dAlso at Università di Bari, Bari, Italy.

^eAlso at Università di Bologna, Bologna, Italy.

^fAlso at Università di Cagliari, Cagliari, Italy.

^gAlso at Università di Ferrara, Ferrara, Italy.

^hAlso at Università di Urbino, Urbino, Italy.

ⁱAlso at Università di Modena e Reggio Emilia, Modena, Italy.

^jAlso at Università di Genova, Genova, Italy.

^kAlso at Università di Milano Bicocca, Milano, Italy.

^lAlso at Università di Roma Tor Vergata, Roma, Italy.

^mAlso at Università di Roma La Sapienza, Roma, Italy.

ⁿAlso at Università della Basilicata, Potenza, Italy.

^oAlso at AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

^pAlso at LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.

^qAlso at Hanoi University of Science, Hanoi, Viet Nam.

^rAlso at Università di Padova, Padova, Italy.

^sAlso at Università di Pisa, Pisa, Italy.

^tAlso at Scuola Normale Superiore, Pisa, Italy.

^uAlso at Università degli Studi di Milano, Milano, Italy.