


Dimensional Analysis of Models and Data Sets: Similarity Solutions and Scaling Analysis



James F. Price

Woods Hole Oceanographic Institution
Woods Hole, Massachusetts, 02543

July 31, 2006

Summary: This essay describes a three-step procedure of dimensional analysis that can be applied to all quantitative models and data sets. In rare but important cases the result of dimensional analysis will be a solution; more often the result is an efficient way to display a large or complex data set.

The first step of an analysis is to define an appropriate physical model, which is nothing more than a list of the dependent variable and all of the independent variables and parameters that are thought to be significant. The premise of dimensional analysis is that a complete equation made from this list of variables will be independent of the choice of units. This leads to the second step, calculation of a null space basis of the corresponding dimensional matrix (a computer code is made available for this calculation). To each vector

of the null space basis there corresponds a nondimensional variable, the number of which is less than the number of dimensional variables. The nondimensional variables are themselves a basis set and in most cases their form is not determined by dimensional analysis alone.

The third and in some respects most interesting step is to choose an optimal form for the basis set. One very useful strategy is to nondimensionalize the dependent variable by a physically motivated 'zero order' solution. When carried to completion, this leads naturally to a scaling analysis in which the nondimensional variables of a model equation are $O(1)$ in some relevant limit. Scaling analysis is often an essential first step in an approximation method. The remaining nondimensional variables can then be formed in ways that define the geometry of the problem or that correspond to the ratios of terms in a model equation, e.g., the Reynolds number or Froude number often arise in models of geophysical fluid dynamics.

Cover page graphic: The tension in the line of a simple pendulum computed by a model for 16 different combinations of length, mass, gravitational acceleration and initial displacement, the angle, ϕ_0 . The upper graph shows the tension as a function of time in dimensional units, and the lower graph is the same data shown in non-dimensional units. The 16 separate solutions collapse to just two, depending upon ϕ_0 and time alone. This kind of data compression illustrates an important advantage to the use of nondimensional variables and dimensional analysis that is described further in Section 4.3.

Contents

1	About dimensional analysis	4
1.1	The goal and the plan	4
1.2	About this essay	5
2	Models of a simple pendulum	5
2.1	A physical model	6
2.2	A mathematical model	6
2.3	Models generally	8
3	An informal dimensional analysis	9
3.1	Invariance to a change of units	9
3.2	Natural units	11
3.3	Extra and omitted variables	12
4	A basis set of nondimensional variables	13
4.1	The mathematical problem	13
4.2	The null space	15
4.3	A basis set for the simple, inviscid pendulum	16

5	The viscous pendulum	18
5.1	A physical model of the viscous pendulum	19
5.2	Drag on a moving sphere	20
5.2.1	Zero order solution	21
5.2.2	The other nondimensional variables: remarks on the Reynolds number	22
5.3	A numerical simulation	23
5.4	An approximate model of the decay rate	25
6	A similarity solution for diffusion in one dimension	27
6.1	Honing the physical model	28
6.2	A similarity solution	28
7	Scaling analysis	31
7.1	A nonlinear projectile problem	31
7.2	Small parameter \rightarrow small term?	34
7.3	Scaling the dependent variable	36
7.4	Approximate and iterated solutions	37
8	Summary and closing remarks	39